SHORT BOWEL SYNDROME

Symptoms Checklist



Short bowel syndrome (SBS) may occur in individuals who have had surgery to remove a portion of their small intestine to treat a damaged small intestine or intestines which have an altered functional ability. The condition may also be congenital.

The signs and symptoms of short bowel syndrome may include:

SIGNS/SYMPTOMS	YES	NO	FREQUENCY	COMMENTS
Bloating				
Cramping				
Dehydration				
Diarrhea				
Fatigue				
Flatulence				
Foul-smelling stool				
Heartburn				
Vomiting				
Weakness				

References/ Resources

Cohran VC, Prozialeck JD, Cole CR. Redefining short bowel syndrome in the 21st century. *Pediatr Res.* 2017 Apr; 81(4):540–549.

Duggan CP, Jaksic T. Pediatric intestinal failure. N Engl J Med. 2017; 377:666-675.

Massironi S, Cavalcoli F, Rausa E, Invernizzi P, Braga M, Vecchi M. Understanding short bowel syndrome: Current status and future perspectives. *Dig Liver Dis.* 2020 Mar; 52(3):253–261.

National Institutes of Health. National Institute of Diabetes and Digestive and Kidney Diseases. Short bowel syndrome. Accessed June 22, 2021. https://www.niddk.nih.gov/health-information/digestive-diseases/short-bowelsyndrome#symptoms.

O'Keefe SJ, Buchman AL, Fishbein TM, Jeejeebhoy KN, Jeppesen PB, Shaffer J. Short bowel syndrome and intestinal failure: consensus definitions and overview. *Clin Gastroenterol Hepatol.* 2006 Jan; 4(1):6–10.

Shakhsheer BA, Warner BW. Short Bowel Syndrome. Curr Treat Options Pediatr. 2019 Dec; 5(4):494-505.